

produced in pairs by the decay of the neutral pions, the cross sections for the processes (1) and (2) would be $(10 \pm 4) \times 10^{-27}$ and $(20 \pm 5) \times 10^{-27}$ cm². The cross section obtained for the charge exchange process is not very sensitive to the angular distribution adopted. It would be $(29 \pm 7) \times 10^{-27}$ cm² for a $\cos^2\theta$ -distribution and $(18 \pm 4) \times 10^{-27}$ cm² for a $\sin^2\theta$ -distribution.

* Research sponsored by the ONR and AEC.

Total Cross Sections of Positive Pions in Hydrogen*

H. L. ANDERSON, E. FERMI, E. A. LONG,† AND D. E. NAGLE
Institute for Nuclear Studies, University of Chicago,
Chicago, Illinois

(Received January 21, 1952)

IN a previous letter,¹ measurements of the total cross sections of negative pions in hydrogen were reported. In the present letter, we report on similar experiments with positive pions.

The experimental method and the equipment used in this measurement was essentially the same as that used in the case of negative pions. The main difference was in the intensity, which for the positives was much less than for the negatives, the more so the higher the energy. This is due to the fact that the positive pions which escape out of the fringing field of the cyclotron magnet are those which are emitted in the backward direction with respect to the proton beam, whereas the negative pions are those emitted in the forward direction. The difficulty of the low intensity was in part compensated by the fact that the cross section for positive pions turned out to be appreciably larger than for negative pions. The results obtained thus far are summarized in Table I.

In Fig. 1 the total cross sections of positive and negative pions are collected. It is quite apparent that the cross section of the positive particles is much larger than that of the negative particles, at least in the energy range from 80 to 150 Mev.

In this letter and in the two preceding ones,^{1,2} the three processes: (1) scattering of positive pions, (2) scattering of negative pions with exchange of charge, and (3) scattering of negative pions without exchange of charge have been investigated. It appears that over a rather wide range of energies, from about 80 to 150 Mev, the cross section for process (1) is the largest, for process (2) is intermediate, and for process (3) is the smallest. Furthermore, the cross sections of both positive and negative pions increase rather rapidly with the energy. Whether the cross sections level off at a high value or go through a maximum, as might be expected if there should be a resonance, is impossible to determine from our present experimental evidence.

Brueckner³ has recently pointed out that the existence of a broad resonance level with spin 3/2 and isotopic spin 3/2 would give an approximate understanding of the ratios of the cross sections for the three processes (1), (2), and (3). We might point out in this connection that the experimental results obtained to date are also compatible with the more general assumption that in the energy interval in question the dominant interaction responsible for the scattering is through one or more intermediate states of isotopic spin 3/2, regardless of the spin. On this assumption, one finds that the ratio of the cross sections for the three

TABLE I. Total cross sections of positive pions in hydrogen.

Energy (Mev)	Cross section (10^{-27} cm ²)
56 ± 8	20 ± 10
82 ± 7	50 ± 13
118 ± 6	91 ± 6
136 ± 6	152 ± 14




FIG. 1. Total cross sections of negative pions in hydrogen (sides of the rectangle represent the error) and positive pions in hydrogen (arms of the cross represent the error). The cross-hatched rectangle is the Columbia result. The black square is the Brookhaven result and does not include the charge exchange contribution.

processes should be (9:2:1), a set of values which is compatible with the experimental observations. It is more difficult, at present, to say anything specific as to the nature of the intermediate state or states. If there were one state of spin 3/2, the angular distribution for all three processes should be of the type $1+3\cos^2\theta$. If the dominant effect were due to a state of spin 1/2, the angular distribution should be isotropic. If states of higher spin or a mixture of several states were involved, more complicated angular distributions would be expected. We intend to explore further the angular distribution in an attempt to decide among the various possibilities.

Besides the angular distribution, another important factor is the energy dependence. Here the theoretical expectation is that, if there is only one dominant intermediate state of spin 3/2 and isotopic spin 3/2, the total cross section of negative pions should at all points be less than $(8/3)\pi\lambda^2$. Apparently, the experimental cross section above 150 Mev is larger than this limit, which indicates that other states contribute appreciably at these energies. Naturally, if a single state were dominant, one could expect that the cross sections would go through a maximum at an energy not far from the energy of the state involved. Unfortunately, we have not been able to push our measurements to sufficiently high energies to check on this point.

Also very interesting is the behavior of the cross sections at low energies. Here the energy dependence should be approximately proportional to the 4th power of the velocity if only states of spin 1/2 and 3/2 and even parity are involved and if the pion is pseudoscalar. The experimental observations in this and other laboratories seem to be compatible with this assumption, but the cross section at low energy is so small that a precise measurement becomes difficult.

* Research sponsored by the ONR and AEC.

† Institute for the Study of Metals, University of Chicago.

¹ Anderson, Fermi, Long, Martin, and Nagle, *Phys. Rev.*, this issue.

² Fermi, Anderson, Lundby, Nagle, and Yodh, preceding Letter, this issue, *Phys. Rev.*

³ K. A. Brueckner (private communication).

Conductivity of Cold-Worked Metals*

D. L. DEXTER

Department of Physics, University of Illinois, Urbana, Illinois

(Received January 17, 1952)

THE change in the electrical conductivity of a metal upon cold-working was first calculated by Koehler,¹ on the assumption that the change is primarily due to the dislocations themselves (rather than associated clusters of vacancies, for example). The scattering potential he used was the difference in